

Topological defect system in $O(n)$ symmetric time-dependent Ginzburg-Landau model

Ying Jiang*

CCAST (World Laboratory), Box 8730, Beijing 100080, P. R. China

Institute of Theoretical Physics, Chinese Academy of Sciences, P. O. Box 2735, Beijing 100080, P. R. China[†]

We present a new generalized topological current in terms of the order parameter field $\vec{\phi}$ to describe the topological defect system in $O(n)$ symmetric time-dependent Ginzburg-Landau model. With the aid of the ϕ -mapping method, the structure of the topological defects and the topological quantization of their topological charges in TDGL model are obtained under the condition that the Jacobian $J(\frac{\phi}{v}) \neq 0$. We show that the topological defects are generated from the zero points of the order parameter field $\vec{\phi}$, and the topological charges of these topological defects are topological quantized in terms of the Hopf indices and Brouwer degrees of ϕ -mapping under the condition. When $J(\frac{\phi}{v}) = 0$, it is shown that there exist the crucial case of branch process. Based on the implicit function theorem and the Taylor expansion, we detail the bifurcation of generalized topological current and find different directions of the bifurcation. The topological defects in TDGL model are found splitting or merging at the degenerate point of field function $\vec{\phi}$ but the total charge of the topological defects is still unchanged.

PACS Numbers: 05.70.Fh, 64.60.Ht, 47.20.Ky, 11.27.+d, 02.40.Pc

I. INTRODUCTION

The importance of the role of time-dependent Ginzburg-Landau (TDGL) model in understanding a variety of problems in physics is clear [1]. The study of the growth kinetics of systems subjected to rapid temperature quenches [2–4] has recently been extended to include systems with more complex order parameter symmetries [5,6]. In particular there has been progress on the study of the n -dimensional vector model with nonconserved Langevin dynamics. An interface description and numerical simulations of a TDGL equation were used to investigate the intra-surface kinetics of phase ordering on toroidal and corrugated surfaces [7]. Two-dimensional XY models with resistively-shunted junction (RSJ) dynamics and TDGL dynamics were simulated and it was verified that the vortex responses can be well described by the Minnhagen phenomenology for both types of dynamics [8].

In recent years, much work has been done on the topological defects in the TDGL model [1,9–11]. In certain cosmological [12] and phase ordering [13] problems key questions involve an understanding of the evolution and correlation among defects like vortices, monopoles, domain walls, etc. In studying such objects in field theory questions arise as to how one can define quantities like the density of topological defects and an associated velocity field. Liu and Mazenko [1,9] have discussed the problem, but unfortunately, for the lack of a powerful method, the topological properties of these systems are not very clear, some important topological information has been lost, also the unified theory of describing the topological properties of all defect system in TDGL model is not established yet.

In our previous work [14], we discussed how one could use the ϕ -mapping topological current theory to study the topological structure of the point defects in phase-ordering systems. We deduced a topological current of point-defect system in terms of the order parameter field in the context of a d -dimensional $O(n)$ symmetric TDGL model, where d is the spatial dimensionality. Using the expression of the topological current, for point defects ($n = d$), the point-defect velocity field was identified, the topological structure and the topological charges of the point-defect system were studied, also the branch process of the point defects was discussed systematically. This analysis will be extended here to the case of arbitrary dimensional defect system in TDGL model where $n = d - k$, i.e. to study the case for all $n \leq d$.

In this paper, in the light of ϕ -mapping topological current theory [15], a useful method which plays a important role in studying the topological invariants [16,17] and the topological structures of physical systems [18,19], we will investigate the topological quantization and the branch process of arbitrary dimensional topological defects in $O(n)$ symmetric TDGL model. We will show that the topological defects are generated from the zero points of the order parameter field $\vec{\phi}$, and their topological charges are quantized in terms of the Hopf indices and Brouwer degrees of

*E-mail: yjiang@itp.ac.cn

[†]mailing address

ϕ -mapping under the condition that the zero points of field $\vec{\phi}$ are regular points. While at the critical points of the order parameter field $\vec{\phi}$, i.e. the limit points and bifurcation points, there exist branch processes, the topological current of defect bifurcates and the topological defects split or merge at such point, this means that the topological defects system is unstable at these points.

This paper is organized as follows. In section 2, we investigate the topological structure of the defect system in TDGL model and give the expression of the defect velocity field. In section 3, we will point out that the topological charges of these defects are the Winding numbers which are determined by the Hopf indices and the Brouwer degrees of the ϕ -mapping. In section 4, we study the branch process of the defect topological current at the limit points, bifurcation points and higher degenerated points systematically by virtue of the ϕ -mapping theory and the implicit function theorem.

II. TOPOLOGICAL STRUCTURE OF THE DEFECT SYSTEM IN TDGL MODEL

We consider a TDGL model for an n -component order parameter $\vec{\phi} = (\phi^1(\vec{r}, t), \dots, \phi^n(\vec{r}, t))$ in d -spatial dimensions ($d - n = k$) governed by the Langevin equation

$$\frac{\partial \vec{\phi}}{\partial t} = \vec{K} \equiv -\Gamma \frac{\delta F}{\delta \vec{\phi}} + \vec{\eta} \quad (1)$$

where Γ is a kinetic coefficient and $\vec{\eta}$ is a thermal noise which is related to Γ by fluctuation-dissipation theorem. F is a Ginzburg-Landau effective free energy assumed to be of the form

$$F = \int d^d r [\frac{1}{2} c (\nabla \vec{\phi})^2 + V(\vec{\phi})] \quad (2)$$

where $c > 0$ and the potential V is assumed to be of the degenerate double-well form. This model is to be supplemented by random, uncorrelated, initial conditions. We assume that there is a rapid temperature quench from a high temperature to zero temperature where the noise $\vec{\eta}$ in (1) can be set to zero. In the scalar case ($n = 1$) such system order through the growth of domains separated by sharp walls. As time evolves these domains coarsen and order grows to progressively longer length scales. In the case of systems with continuous symmetry ($n > 1$) the disordering elements [20,21] will depend on n and spatial dimensionality d . Thus, for example, for $n = d$ one has point defects (vortices and monopoles), for $n = d - 1$ one has vortex line or string-like defects while for $n = d - k$ one has k -dimensional topological defect objects. For $n > d$ there are no stable singular topological objects.

As is well known, the n -component order parameter field $\vec{\phi}(\vec{r}, t)$ determines the defect properties of the system, and it can be looked upon as a smooth mapping between the $(d + 1)$ -dimensional space-time X and the n -dimensional Euclidean space R^n as $\phi : X \rightarrow R^n$. By analogy with the discussion in our previous work [14,15,19,22], from this ϕ -mapping, one can deduce a topological tensor current as

$$j^{\mu_0 \mu_1 \dots \mu_k} = \frac{1}{A(S^{n-1})(n-1)!} \epsilon^{\mu_0 \mu_1 \dots \mu_k \mu_{k+1} \dots \mu_d} \epsilon_{a_1 \dots a_n} \partial_{\mu_{k+1}} n^{a_1} \partial_{\mu_{k+2}} n^{a_2} \dots \partial_{\mu_d} n^{a_n},$$

$$\mu_0, \dots, \mu_d = 0, 1, \dots, d, \quad a_1, \dots, a_n = 1, \dots, n \quad (3)$$

to describe the k -dimensional topological defects in TDGL model, and the density tensor of the topological defect system is defined as $\rho^{\mu_1 \dots \mu_k} = j^{0 \mu_1 \dots \mu_k}$. In this expression, ∂_μ stands for $\partial/\partial x^\mu$, $A(S^{n-1}) = 2\pi^{n/2}/\Gamma(n/2)$ is the area of $(n - 1)$ -dimensional unit sphere S^{n-1} and $n^a(x)$ is the direction field of the n -component order parameter field $\vec{\phi}$

$$n^a(x) = \frac{\phi^a(x)}{\|\phi(x)\|}, \quad \|\phi(x)\| = \sqrt{\phi^a(x)\phi^a(x)} \quad (4)$$

with $n^a(x)n^a(x) = 1$. It is obviously that $n^a(x)$ is a section of the sphere bundle $S(X)$ [15] and it can be looked upon as a map of X onto a $(n - 1)$ -dimensional unit sphere S^{n-1} in order parameter space. Clearly, the zero points of the order parameter field $\vec{\phi}(x)$ are just the singular points of the unit vector $n^a(x)$. It is easy to see that $j^{\mu_0 \dots \mu_k}$ are completely antisymmetric, and from the formulas above, we conclude that there exists a conservative equation of the topological tensor current in (3)

$$\partial_{\mu_i} j^{\mu_0 \dots \mu_k} = 0, \quad i = 0, \dots, k,$$

and from which we have

$$\partial_t \rho^{\mu_1 \cdots \mu_k} + \nabla_\mu j^{\mu \mu_1 \cdots \mu_k} = 0$$

which is just the continuity equation satisfied by $\rho^{\mu_1 \cdots \mu_k}$.

In the following, we will investigate the intrinsic structure of the generalized topological current $j^{\mu_0 \mu_1 \cdots \mu_k}$ by making use of the ϕ -mapping method. From (4), we have

$$\partial_\mu n^a = \frac{1}{\|\phi\|} \partial_\mu \phi^a + \phi^a \partial_\mu \left(\frac{1}{\|\phi\|} \right), \quad \frac{\partial}{\partial \phi^a} \left(\frac{1}{\|\phi\|} \right) = -\frac{\phi^a}{\|\phi\|^3}$$

which should be looked upon as generalized functions [23]. Due to these expressions the generalized topological current (3) can be rewritten as

$$j^{\mu_0 \mu_1 \cdots \mu_k} = C_n \epsilon^{\mu_0 \mu_1 \cdots \mu_k \mu_{k+1} \cdots \mu_d} \epsilon_{a_1 \cdots a_n} \cdot \partial_{\mu_{k+1}} \phi^{a_1} \cdots \partial_{\mu_d} \phi^{a_n} \frac{\partial}{\partial \phi^{a_1}} \frac{\partial}{\partial \phi^{a_1}} (G_n(\|\phi\|)), \quad (5)$$

where C_n is a constant

$$C_n = \begin{cases} -\frac{1}{A(S^{n-1})(n-2)(n-1)!}, & n > 2 \\ \frac{1}{2\pi}, & n = 2 \end{cases},$$

and $G_n(\|\phi\|)$ is a Green function

$$G_n(\|\phi\|) = \begin{cases} \frac{1}{\|\phi\|^{n-2}}, & n > 2 \\ \ln \|\phi\|, & n = 2 \end{cases}.$$

Defining general Jacobians $J^{\mu_0 \mu_1 \cdots \mu_k}(\frac{\phi}{x})$ as following

$$\epsilon^{a_1 \cdots a_n} J^{\mu_0 \mu_1 \cdots \mu_k}(\frac{\phi}{x}) = \epsilon^{\mu_0 \mu_1 \cdots \mu_k \mu_{k+1} \cdots \mu_d} \partial_{\mu_{k+1}} \phi^{a_1} \partial_{\mu_{k+2}} \phi^{a_2} \cdots \partial_{\mu_d} \phi^{a_n}$$

and by making use of the n -dimensional Laplacian Green function relation in ϕ -space [15]

$$\Delta_\phi (G_n(\|\phi\|)) = -\frac{4\pi^{n/2}}{\Gamma(\frac{n}{2} - 1)} \delta(\vec{\phi})$$

where $\Delta_\phi = (\frac{\partial^2}{\partial \phi^a \partial \phi^a})$ is the n -dimensional Laplacian operator in ϕ -space, we do obtain the δ -function structure of the defect topological current rigorously

$$j^{\mu_0 \mu_1 \cdots \mu_k} = \delta(\vec{\phi}) J^{\mu_0 \mu_1 \cdots \mu_k}(\frac{\phi}{x}). \quad (6)$$

This expression involves the total defect information of the system in TDGL model and it indicates that all the defects are located at the zero points of the order parameter field $\vec{\phi}(x)$. It must be pointed out that, comparing to similar expressions in other papers, the results in (6) is gotten theoretically in a natural way. From this expression, the density tensor of $j^{\mu_0 \mu_1 \cdots \mu_k}$ is also changed into a compact form

$$\rho^{\mu_1 \cdots \mu_k} = j^{0 \mu_1 \cdots \mu_k} = \delta(\vec{\phi}) D^{\mu_1 \cdots \mu_k}(\frac{\phi}{x}),$$

where $D^{\mu_1 \cdots \mu_k}(\frac{\phi}{x}) = J^{0 \mu_1 \cdots \mu_k}(\frac{\phi}{x})$. We find that $j^{\mu_0 \mu_1 \cdots \mu_k} \neq 0$, $\rho^{\mu_1 \cdots \mu_k} \neq 0$ only when $\vec{\phi} = 0$, which is just the singularity of $j^{\mu_0 \mu_1 \cdots \mu_k}$. In detail, the Kernel of the ϕ -mapping is the singularities of the topological tensor current $j^{\mu_0 \mu_1 \cdots \mu_k}$ in X , i.e. the inner structure of the topological tensor current is labeled by the zeroes of ϕ -mapping. We think that this is the essential of the topological tensor current theory and ϕ -mapping is the key to study this theory.

From the above discussions, we see that the kernel of ϕ -mapping plays an important role in the topological tensor current theory, so we are focused on the zero points of $\vec{\phi}$ and will search for the solutions of the equations $\phi^a(x) = 0$ ($a = 1, \dots, n$) by means of the implicit function theorem. These points are topological singularities in the orientation of the order parameter field $\vec{\phi}(x)$.

Suppose that the vector field $\vec{\phi}(x)$ possesses l zeroes, according to the implicit function theorem [24], when the zeroes are regular points of ϕ -mapping at which the rank of the Jacobian matrix $[\partial_\mu \phi^a]$ is n , the solutions of $\vec{\phi} = 0$ can be expressed parameterizedly by

$$x^\mu = z_i^\mu(t, u^1, \dots, u^k), \quad i = 1, \dots, l, \quad (7)$$

where the subscript i represents the i -th solution and the parameters u^I ($I = 1, \dots, k$), combining with the time parameter t , span a $(k+1)$ -dimensional subspace which is called the i -th singular subspace N_i in the space-time X corresponding to the ϕ -mapping. These singular subspaces N_i are just the world volumes of the topological defects, the parameters u^I play the role of the spatial parameters of the topological defects. For each singular subspace N_i , we can define a normal subspace M_i in X which is spanned by the parameters v^A ($A = 1, \dots, n$), and the intersection point of M_i and N_i is denoted by p_i which can be expressed parameterizedly by $v^A = p_i^A$. In fact, in the words of differential topology, M_i is transversal to N_i at the point p_i . By virtue of the implicit function theorem at the regular point p_i , it should be held true that the Jacobian matrices $J(\frac{\phi}{v})$ satisfies $J(\frac{\phi}{v}) = \frac{D(\phi^1, \dots, \phi^n)}{D(v^1, \dots, v^n)} \neq 0$.

On the other hand, putting the solutions (7) into $\vec{\phi}(x)$, we have $\vec{\phi}(z_i) \equiv 0$, from this expression, and since we expect the instantaneous velocity to be orthogonal to the local orientation of the topological defects, we can define the velocity via

$$J^{\mu\mu_1 \dots \mu_k} = v^{[\mu} D^{\mu_1 \dots \mu_k]}, \quad (8)$$

that is, the topological current and the density tensor should satisfy

$$j^{\mu\mu_1 \dots \mu_k} = v^{[\mu} \rho^{\mu_1 \dots \mu_k]}. \quad (9)$$

This expression for the velocity can be used to find the defect velocity distribution in the case of phase-ordering kinetics for a non-conserved order parameter [1,9,14]. It is very useful because it avoid the problem of having to specify the positions of the topological defects explicitly. The positions are implicitly determined by the zeros of the order parameter field. The general expression with $J^{\mu\mu_1 \dots \mu_k}(\phi/x)$ should be useful in looking at the motion of the topological defects in TDGL model in the presence of external fields beyond a growth kinetics context [1,9].

If we restrict the discussion to the simplest case of point defects ($n = d$), the corresponding topological current is reduced to $j^\mu = \delta(\phi) J^\mu(\phi/x)$, and Eq.(9) can be put into a conventional form $j^\mu = v^\mu \rho$ with v^μ taking the expression of $v^\mu = J^\mu(\phi/x)/D(\phi/x)$ [14]. Also, the string in TDGL model, which was discussed by Mazenko [25], is a special case for $n = d - 1$ in our theory.

III. TOPOLOGICAL QUANTIZATION OF THE DEFECT CHARGES IN TDGL MODEL

In the following, we will investigate the topological charges of the topological defects and their quantization. Let Σ_i be a neighborhood of p_i on M_i with boundary $\partial\Sigma_i$ satisfying $p_i \notin \partial\Sigma_i$, $\Sigma_i \cap \Sigma_j = \emptyset$. Then the generalized winding number W_i of $n^a(x)$ at p_i [26] can be defined by the Gauss map $n : \partial\Sigma_i \rightarrow S^{n-1}$

$$W_i = \frac{1}{A(S^{n-1})(n-1)!} \int_{\partial\Sigma_i} n^* (\epsilon_{a_1 \dots a_n} n^{a_1} dn^{a_2} \wedge \dots \wedge dn^{a_n}) \quad (10)$$

where n^* denotes the pull back of map n . The generalized winding numbers is a topological invariant and is also called the degree of Gauss map [27]. It means that, when the point v^A covers $\partial\Sigma_i$ once, the unit vector n^a will cover a region $n[\partial\Sigma_i]$ whose area is W_i times of $A(S^{n-1})$, i.e. the unit vector n^a will cover the unit sphere S^{n-1} for W_i times. Using the Stokes' theorem in exterior differential form and duplicating the above process, we get the compact form of W_i

$$W_i = \int_{\Sigma_i} \delta(\vec{\phi}) J(\frac{\phi}{v}) d^n v. \quad (11)$$

By analogy with the procedure of deducing $\delta(f(x))$, since

$$\delta(\vec{\phi}) = \begin{cases} +\infty, & \text{for } \vec{\phi}(x) = 0 \\ 0, & \text{for } \vec{\phi}(x) \neq 0 \end{cases} = \begin{cases} +\infty, & \text{for } x \in N_i \\ 0, & \text{for } x \notin N_i \end{cases}, \quad (12)$$

we can expand the δ -function $\delta(\vec{\phi})$ as

$$\delta(\vec{\phi}) = \sum_{i=1}^l c_i \delta(N_i), \quad (13)$$

where the coefficients c_i must be positive, i.e. $c_i = |c_i|$. $\delta(N_i)$ is the δ -function in space-time X on a submanifold N_i [23]

$$\delta(N_i) = \int_{N_i} \delta^n(\vec{x} - \vec{z}_i(t, u^1, \dots, u^k)) dt d^k u. \quad (14)$$

Substituting (13) into (11), and calculating the integral, we get the expression of c_i

$$c_i = \frac{\beta_i}{|J(\frac{\phi}{v})_{p_i}|} = \frac{\beta_i \eta_i}{J(\frac{\phi}{v})_{p_i}}, \quad (15)$$

where $\beta_i = |W_i|$ is a positive integer called the Hopf index [27] of ϕ -mapping on M_i , it means that when the point v covers the neighborhood of the zero point p_i once, the function $\vec{\phi}$ covers the corresponding region in $\vec{\phi}$ -space β_i times, and $\eta_i = \text{sign} J(\frac{\phi}{v})_{p_i} = \pm 1$ is the Brouwer degree of ϕ -mapping [27]. Substituting this expression of c_i and (13) into (6), we gain the total expansion of the topological current

$$j^{\mu_0 \mu_1 \dots \mu_k} = \sum_{i=1}^l \frac{\beta_i \eta_i}{J(\frac{\phi}{v})_{p_i}} \delta(N_i) J^{\mu_0 \mu_1 \dots \mu_k}(\frac{\phi}{x}).$$

or in terms of parameters $y^{A'} = (t, v^1, \dots, v^n, u^1, \dots, u^k)$

$$j^{A'_0 A'_1 \dots A'_k} = \sum_{i=1}^l \frac{\beta_i \eta_i}{J(\frac{\phi}{v})_{p_i}} \delta(N_i) J^{A'_0 A'_1 \dots A'_k}(\frac{\phi}{y}). \quad (16)$$

From the above equation, we conclude that the inner structure of $j^{\mu_0 \mu_1 \dots \mu_k}$ or $j^{A'_0 A'_1 \dots A'_k}$ is labeled by the total expansion of $\delta(\vec{\phi})$, and it just represents l (k)-dimensional topological defects with topological charges $g_i = \beta_i \eta_i$ moving in the $(d+1)$ -dimensional space-time X . The $(k+1)$ -dimensional singular subspaces N_i ($i = 1, \dots, l$) are their world sheets in the space-time. Mazenko [1,9] and Halperin [28] also got similar results for the case of point-like defects and line defects, but unfortunately, they did not consider the case $\beta_i \neq 1$. In fact, what they lost sight of is just the most important topological information for the charge of topological defects. In detail, the Hopf indices β_i characterize the absolute values of the topological charges of these defects and the Brouwer degrees $\eta_i = +1$ correspond to defects while $\eta_i = -1$ to antidefects. Furthermore, they did not discuss what will happen when η_i is indefinite, which we will study in detail in section 4.

IV. THE BRANCH PROCESS OF THE TOPOLOGICAL CURRENT

With the discussion mentioned above, we know that the results in the above section are obtained straightly from the topological view point under the condition $J(\phi/v)|_{p_i} \neq 0$, i.e. at the regular points of the order parameter field $\vec{\phi}$. When the condition fails, i.e. the Brouwer degree η_i are indefinite, there should exist some kind of branch processes in the topological current of the topological defect system in TDGL model. In what follows, we will study the case when $J(\phi/v)|_{p_i} = 0$. It often happens when the zero points of field $\vec{\phi}$ include some branch points, which lead to the bifurcation of the topological current.

In this section, we will discuss the branch processes of these topological defects. In order to simplify our study, let the spatial parameters u^I be fixed, i.e. to choose a fixed point on the topological defect. In this case, the Jacobian matrices $J^{A'_0 A'_1 \dots A'_k}(\frac{\phi}{y})$ are reduced to

$$J^{A I_1 \dots I_k}(\frac{\phi}{y}) \equiv J^A(\frac{\phi}{y}), \quad J^{A B I_1 \dots I_{k-1}}(\frac{\phi}{y}) = 0, \quad J^{(n+1) \dots d}(\frac{\phi}{y}) = J(\frac{\phi}{v}),$$

$$A, B = 0, 1, \dots, n, \quad I_j = n+1, \dots, d, \quad (17)$$

for $y^A = v^A$ ($A \leq n$), $y^0 = t$, $y^{n+I} = u^I$ ($I \geq 1$). The branch points are determined by the $n + 1$ equations

$$\phi^a(t, v^1, \dots, v^n, \vec{u}) = 0, \quad a = 1, \dots, n \quad (18)$$

and

$$\phi^{n+1}(t, v^1, \dots, v^n, \vec{u}) \equiv J\left(\frac{\phi}{v}\right) = 0 \quad (19)$$

for the fixed $\vec{u}$. and they are denoted as (t^*, p_i) . In ϕ -mapping theory usually there are two kinds of branch points, namely the limit points and bifurcation points [29], satisfying

$$J^1\left(\frac{\phi}{y}\right)|_{(t^*, p_i)} \neq 0 \quad (20)$$

and

$$J^1\left(\frac{\phi}{y}\right)|_{(t^*, p_i)} = 0, \quad (21)$$

respectively. In the following, we assume that the branch points (t^*, p_i) of ϕ -mapping have been found.

A. The branch process at the limit point

We first discuss the branch process at the limit point satisfying the condition (20). In order to use the theorem of implicit function to study the branch process of topological defects at the limit point, we use the Jacobian $J^1(\frac{\phi}{y})$ instead of $J(\frac{\phi}{v})$ to discuss the problem. In fact, this means that we have replaced the parameter t by v^1 . Then, taking account of the condition(20) and using the implicit function theorem, we have an unique solution of the equations (18) in the neighborhood of the limit point (t^*, p_i)

$$t = t(v^1, \vec{u}), \quad v^i = v^i(v^1, \vec{u}), \quad i = 2, 3, \dots, n \quad (22)$$

with $t^* = t(p_i^1, \vec{u})$. In order to show the behavior of the defects at the limit points, we will investigate the Taylor expansion of (22) in the neighborhood of (t^*, p_i) . In the present case, from (20) and (19), we get

$$\frac{dv^1}{dt}|_{(t^*, p_i)} = \frac{J^1(\frac{\phi}{y})}{J(\frac{\phi}{v})}|_{(t^*, p_i)} = \infty,$$

i.e.

$$\frac{dt}{dv^1}|_{(t^*, p_i)} = 0.$$

Therefore, the Taylor expansion of (22) at the point (t^*, p_i) gives

$$t - t^* = \frac{1}{2} \frac{d^2 t}{(dv^1)^2}|_{(t^*, p_i)} (v^1 - p_i^1)^2 \quad (23)$$

which is a parabola in the v^1 — t plane. From (23), we can obtain the two solutions $v_{(1)}^1(t, \vec{u})$ and $v_{(2)}^1(t, \vec{u})$, which give the branch solutions of the system (18) at the limit point. If $\frac{d^2 t}{(dv^1)^2}|_{(t^*, p_i)} > 0$, we have the branch solutions for $t > t^*$ (Fig 1(a)), otherwise, we have the branch solutions for $t < t^*$ (Fig 1(b)). The former is related to the creation of defect and antidefect in pair at the limit points, and the latter to the annihilation of the topological defects, since the topological current of topological defects is identically conserved, the topological quantum numbers of these two generated topological defects must be opposite at the limit point, i.e. $\beta_1 \eta_1 + \beta_2 \eta_2 = 0$.

B. The branch process at the bifurcation point

In the following, let us consider the case (21), in which the restrictions of the system (18) at the bifurcation point (t^*, p_i) are

$$J\left(\frac{\phi}{v}\right)|_{(t^*, p_i)} = 0, \quad J^1\left(\frac{\phi}{y}\right)|_{(t^*, p_i)} = 0. \quad (24)$$

These two restrictive conditions will lead to an important fact that the dependency relationship between t and v^1 is not unique in the neighborhood of the bifurcation point (t^*, p_i) . In fact, we have

$$\frac{dv^1}{dt}|_{(t^*, p_i)} = \frac{J^1\left(\frac{\phi}{y}\right)|_{(t^*, p_i)}}{J\left(\frac{\phi}{v}\right)|_{(t^*, p_i)}} \quad (25)$$

which under the restraint (24) directly shows that the tangential direction of the integral curve of equation (25) is indefinite at the point (t^*, p_i) . Hence, (25) does not satisfy the conditions of the existence and uniqueness theorem of the solution of a differential equation. This is why the very point (t^*, z_i) is called the bifurcation point of the system (18).

As we have mentioned above, at the bifurcation point (t^*, p_i) , the rank of the Jacobian matrix $[\frac{\partial \phi}{\partial v}]$ is smaller than n . For the aim of searching for the different directions of all branch curves at the bifurcation point, we firstly consider the rank of the Jacobian matrix $[\frac{\partial \phi}{\partial v}]$ is $n - 1$. The case of a more smaller rank will be discussed in next subsection. Let $J_1(\frac{\phi}{v}) = [\phi_A^a]$ ($a = 1, \dots, n - 1$; $A = 2, \dots, n$) be one of the $(n - 1) \times (n - 1)$ submatrix of the Jacobian matrix $[\frac{\partial \phi}{\partial v}]$ with $\det J_1(\frac{\phi}{v}) \neq 0$ at the point (t^*, p_i) (otherwise, we have to rearrange the equations of (18)), where ϕ_A^a stands for $(\partial \phi^a / \partial v^A)$. By means of the implicit function theorem we obtain one and only one functional relationship in the neighborhood of the bifurcation point (t^*, p_i)

$$v^A = f^A(v^1, t, \vec{u}), \quad A = 2, 3, \dots, n. \quad (26)$$

We denote the partial derivatives as $f_1^A = \frac{\partial v^A}{\partial v^1}$, $f_t^A = \frac{\partial v^A}{\partial t}$, $f_{11}^A = \frac{\partial^2 v^A}{(\partial v^1)^2}$, $f_{1t}^A = \frac{\partial^2 v^A}{\partial v^1 \partial t}$, $f_{tt}^A = \frac{\partial^2 v^A}{\partial t^2}$. From (18) and (26), we have for $a = 1, \dots, n - 1$

$$\phi^a = \phi^a(v^1, f^2(v^1, t, \vec{u}), \dots, f^n(v^1, t, \vec{u}), t, \vec{u}) \equiv 0$$

which gives

$$\sum_{A=2}^n \frac{\partial \phi^a}{\partial v^A} f_1^A = -\frac{\partial \phi^a}{\partial v^1}, \quad a = 1, \dots, n - 1 \quad (27)$$

$$\sum_{A=2}^n \frac{\partial \phi^a}{\partial v^A} f_t^A = -\frac{\partial \phi^a}{\partial t}, \quad a = 1, \dots, n - 1. \quad (28)$$

By differentiating (27) and (28) with respect to v^1 and t , and applying the Gaussian elimination method, we can find the second order derivatives f_{11}^A , f_{1t}^A and f_{tt}^A . The above discussions do not matter to the last component $\phi^n(v^1, \dots, v^n, t, \vec{u})$. In order to find the different values of dv^1/dt at the bifurcation point, let us investigate the Taylor expansion of $\phi^n(v^1, \dots, v^n, t, \vec{u})$ in the neighborhood of (t^*, p_i) . Substituting (26) into $\phi^n(v^1, \dots, v^n, t, \vec{u})$, we get the function of two variables v^1 and t

$$F(t, v^1, \vec{u}) = \phi^n(v^1, f^2(v^1, t, \vec{u}), \dots, f^n(v^1, t, \vec{u}), t, \vec{u}) \quad (29)$$

which according to (18) must vanish at the bifurcation point

$$F(t^*, p_i) = 0. \quad (30)$$

From (29), we can calculate the first order partial derivatives of $F(t, v^1, \vec{u})$ with respect to v^1 and t respectively at the bifurcation point (t^*, p_i)

$$\frac{\partial F}{\partial v^1} = \phi_1^n + \sum_{A=2}^n \phi_A^n f_1^A, \quad \frac{\partial F}{\partial t} = \phi_t^n + \sum_{A=2}^n \phi_A^n f_t^A. \quad (31)$$

By making use of (27) and (28), with the Cramer's rule, the first equation of (24) is expressed as

$$\frac{\partial F}{\partial v^1} \det J_1\left(\frac{\phi}{v}\right)|_{(t^*, p_i)} = 0.$$

Since $\det J_1\left(\frac{\phi}{v}\right)|_{(t^*, p_i)} \neq 0$, the above equation leads to

$$\frac{\partial F}{\partial v^1}|_{(t^*, p_i)} = 0. \quad (32)$$

With the same reasons, we can prove that

$$\frac{\partial F}{\partial t}|_{(t^*, p_i)} = 0. \quad (33)$$

The second order partial derivatives of the function $F(t, v^1, \vec{u})$ are easily to find out from (31) which at (t^*, p_i) are denoted by

$$A = \frac{\partial^2 F}{(\partial v^1)^2}|_{(t^*, p_i)}, \quad B = \frac{\partial^2 F}{\partial v^1 \partial t}|_{(t^*, p_i)}, \quad C = \frac{\partial^2 F}{\partial t^2}|_{(t^*, p_i)}. \quad (34)$$

Then, by virtue of (30), (32), (33) and (34), the Taylor expansion of $F(t, v^1, \vec{u})$ in the neighborhood of the bifurcation point (t^*, p_i) gives

$$A(v^1 - p_i^1)^2 + 2B(v^1 - p_i^1)(t - t^*) + C(t - t^*)^2 = 0. \quad (35)$$

Dividing (35) by $(v^1 - p_i^1)^2$ or $(t - t^*)^2$, and taking the limit $t \rightarrow t^*$ as well as $v^1 \rightarrow p_i^1$ respectively, we get two equations

$$A\left(\frac{dv^1}{dt}\right)^2 + 2B\frac{dv^1}{dt} + C = 0. \quad (36)$$

and

$$C\left(\frac{dt}{dv^1}\right)^2 + 2B\frac{dt}{dv^1} + A = 0. \quad (37)$$

So we get the different directions of the branch curves at the bifurcation point from the solutions of (36) or (37). There are four possible cases:

Firstly, $A \neq 0$, $\Delta = 4(B^2 - AC) > 0$, from Eq. (36) we get two different solutions: $dv^1/dt|_{1,2} = (-B \pm \sqrt{B^2 - AC})/A$, which is shown in Fig. 2, where two topological defects meet and then depart at the bifurcation point. Secondly, $A \neq 0$, $\Delta = 4(B^2 - AC) = 0$, there is only one solution: $dv^1/dt = -B/A$, which includes three important cases: (a) two topological defects tangentially collide at the bifurcation point (Fig 3(a)); (b) two topological defects merge into one topological defect at the bifurcation point (Fig 3(b)); (c) one topological defect splits into two topological defects at the bifurcation point (Fig 3(c)). Thirdly, $A = 0$, $C \neq 0$, $\Delta = 4(B^2 - AC) > 0$, from Eq. (37) we have $dt/dv^1 = 0$ and $-2B/C$. There are two important cases: (i) One topological defect splits into three topological defects at the bifurcation point (Fig 4(a)); (ii) Three topological defects merge into one at the bifurcation point (Fig 4(b)). Finally, $A = C = 0$, Eqs. (36) and (37) give respectively $dv^1/dt = 0$ and $dt/dv^1 = 0$. This case is obvious as in Fig. 5, which is similar to the third situation.

In order to determine the branches directions of the remainder variables, we will use the relations simply

$$dv^A = f_1^A dv^1 + f_t^A dt, \quad A = 2, 3, \dots, n$$

where the partial derivative coefficients f_1^A and f_t^A have given in (27) and (28). Then, respectively

$$\frac{dv^A}{dv^1} = f_1^A + f_t^A \frac{dt}{dv^1}$$

or

$$\frac{dv^A}{dt} = f_1^A \frac{dv^1}{dt} + f_t^A. \quad (38)$$

where partial derivative coefficients f_1^A and f_t^A are given by (27) and (28). From this relations we find that the values of dv^A/dt at the bifurcation point (t^*, z_i) are also possibly different because (37) may give different values of dv^1/dt .

C. The branch process at the higher degenerated point

In the following, let us discuss the branch process at a higher degenerated point. In the above subsection, we have analyzed the case that the rank of the Jacobian matrix $[\partial\phi/\partial v]$ of the equation (19) is $n - 1$. In this section, we consider the case that the rank of the Jacobian matrix is $n - 2$ (for the case that the rank of the matrix $[\partial\phi/\partial v]$ is lower than $n - 2$, the discussion is in the same way). Let the $(n - 2) \times (n - 2)$ submatrix $J_2(\frac{\phi}{v})$ of the Jacobian matrix $[\partial\phi/\partial v]$ be

$$J_2\left(\frac{\phi}{v}\right) = \begin{pmatrix} \phi_3^1 & \phi_4^1 & \cdots & \phi_n^1 \\ \phi_3^2 & \phi_4^2 & \cdots & \phi_n^2 \\ \vdots & \vdots & \ddots & \vdots \\ \phi_3^{n-2} & \phi_4^{n-2} & \cdots & \phi_n^{n-2} \end{pmatrix}$$

and suppose that $\det J_2(\frac{\phi}{v})|_{(t^*, p_i)} \neq 0$. With the same reasons of obtaining (26), we can have the function relations

$$v^A = f^A(v^1, v^2, t, \vec{u}), \quad A = 3, 4, \dots, n. \quad (39)$$

For the partial derivatives f_1^A , f_2^A and f_t^A , we can easily derive the system similar to the equations (27) and (28), in which the three terms at the right hand of can be figured out at the same time. In order to determine the 2-order partial derivatives f_{11}^A , f_{12}^A , f_{1t}^A , f_{22}^A , f_{2t}^A and f_{tt}^A , we can use the method similar to the above mentioned. Substituting the relations (39) into the last two equations of the system (18), we have the following two equations with respect to the arguments v^1 , v^2 , t , $\vec{u}$

$$\begin{cases} F_1(v^1, v^2, t, \vec{u}) = \phi^{n-1}(v^1, v^2, f^3(v^1, v^2, t, \vec{u}), \dots, f^n(v^1, v^2, t, \vec{u}), t, \vec{u}) = 0 \\ F_2(v^1, v^2, t, \vec{u}) = \phi^n(v^1, v^2, f^3(v^1, v^2, t, \vec{u}), \dots, f^n(v^1, v^2, t, \vec{u}), t, \vec{u}) = 0. \end{cases} \quad (40)$$

Calculating the partial derivatives of the function F_1 and F_2 with respect to v^1 , v^2 and t , taking notice of (39) and using six similar expressions to (32) and (33), i.e.

$$\frac{\partial F_j}{\partial v^1} |_{(t^*, p_i)} = 0, \quad \frac{\partial F_j}{\partial v^2} |_{(t^*, p_i)} = 0, \quad \frac{\partial F_j}{\partial t} |_{(t^*, p_i)} = 0, \quad j = 1, 2, \quad (41)$$

we have the following forms of Taylor expressions of F_1 and F_2 in the neighborhood of (t^*, p_i)

$$F_j(v^1, v^2, t, \vec{u}) \approx A_{j1}(v^1 - p_i^1)^2 + A_{j2}(v^1 - p_i^1)(v^2 - p_i^2) + A_{j3}(v^1 - p_i^1) \cdot (t - t^*) + A_{j4}(v^2 - p_i^2)^2 + A_{j5}(v^2 - p_i^2)(t - t^*) + A_{j6}(t - t^*)^2 = 0$$

$$j = 1, 2. \quad (42)$$

In the case of $A_{j1} \neq 0, A_{j4} \neq 0$, by dividing (42) by $(t - t^*)^2$ and taking the limit $t \rightarrow t^*$, we obtain two quadratic equations of $\frac{dv^1}{dt}$ and $\frac{dv^2}{dt}$

$$A_{j1}\left(\frac{dv^1}{dt}\right)^2 + A_{j2}\frac{dv^1}{dt}\frac{dv^2}{dt} + A_{j3}\frac{dv^1}{dt} + A_{j4}\left(\frac{dv^2}{dt}\right)^2 + A_{j5}\frac{dv^2}{dt} + A_{j6} = 0 \quad (43)$$

$$j = 1, 2.$$

Eliminating the variable dv^1/dt , we obtain a equation of dv^2/dt in the form of a determinant

$$\begin{vmatrix} A_{11} & A_{12}Q + A_{23} & A_{14}Q^2 + A_{15}Q + A_{16} & 0 \\ 0 & A_{11} & A_{12}Q + A_{13} & A_{14}Q^2 + A_{15}Q + A_{16} \\ A_{21} & A_{22}Q + A_{23} & A_{24}Q^2 + A_{25}Q + A_{26} & 0 \\ 0 & A_{21} & A_{22}Q + A_{23} & A_{24}Q^2 + A_{25}Q + A_{26} \end{vmatrix} = 0 \quad (44)$$

where $Q = dv^2/dt$, which is a 4th order equation of dv^2/dt

$$a_0\left(\frac{dv^2}{dt}\right)^4 + a_1\left(\frac{dv^2}{dt}\right)^3 + a_2\left(\frac{dv^2}{dt}\right)^2 + a_3\left(\frac{dv^2}{dt}\right) + a_4 = 0. \quad (45)$$

Therefore we get different directions at the higher degenerated point corresponding to different branch curves. The number of different branch curves is four at most. If the degree of degeneracy of the matrix $[\frac{\partial\phi}{\partial v}]$ is more higher, i.e. the rank of the matrix $[\frac{\partial\phi}{\partial v}]$ is more lower than the present $(n-2)$ case, the procedure of deduction will be more complicate. In general supposing the rank of the matrix $[\frac{\partial\phi}{\partial x}]$ be $(n-s)$, the number of the possible different directions of the branch curves is 2^s at most.

At the end of this section, we conclude that there exist crucial cases of branch processes in our theory of topological defect system in TDGL model. This means that a topological defect, at the bifurcation point, may split into several (for instance m) topological defects along different branch curves with different charges. Since the topological current is a conserved current, the total quantum number of the splitting topological defects must precisely equal to the topological charge of the original defect i.e.

$$\sum_{j=1}^m \beta_{i_j} \eta_{i_j} = \beta_i \eta_i$$

for fixed i . This can be looked upon as the topological reason of the defect splitting. Here we should point out that such splitting is a stochastic process, the sole restriction of this process is just the conservation of the topological charge of the topological defects during this splitting process. Of course, the topological charge of each splitting defects is an integer.

ACKNOWLEDGMENT

The author gratefully acknowledges the support of K. C. Wong Education Foundation, Hong Kong. The author is also gratefully indebted to Prof. Y. S. Duan for his warm-hearted helps and useful discussion.

-
- [1] G. Mazenko, *Phys. Rev. Lett.* **78**, 401 (1997); *Phys. Rev.* **E 59**, 1574 (1999)
 - [2] C. Domb and J. L. Lebowitz, *Phase transitions and critical phenomena*, Vol. 8 (Academic, New York, 1983); H. Furukawa, *Adv. Phys.* **34**, 703 (1985); K. Binder, *Rep. Prog. Phys.* **50**, 783 (1987)
 - [3] K. Kawasaki, M. C. Yalabik and J. D. Gunton, *Phys. Rev. A* **17**, 455 (1978); T. Ohta, D. Jasnow and K. Kawasaki, *Phys. Rev. Lett* **49**, 1223 (1982); A. J. Bray, *Phys. Rev. Lett* **62**, 2841 (1989); *Phys. Rev. B* **41**, 6724 (1990)
 - [4] G. Mazenko, O. Valls, and M. Zannetti, *Phys. Rev. B* **38**, 520 (1988); G. Mazenko, *Phys. Rev. Lett.* **63**, 1605 (1989); *Phys. Rev. B* **42**, 4487 (1990)
 - [5] H. Toyoki, *Phys. Rev. B* **45**, 1965 (1992); A. J. Bray and K. Humayun, *J. Phys. A* **25**, 2191 (1992); F. Liu, M. Monello and N. Goldenfeld, *Phys. Rev. Lett.* **66**, 3071 (1991); H. Frahm, S. Ullah and A. Dorsey, *Phys. Rev. Lett.* **66**, 3067 (1991)
 - [6] M. Zannetti and G. Mazenko, *Phys. Rev. B* **35**, 3066 (1987); A. Bray and S. Puri, *Phys. Rev. Lett.* **67**, 2670 (1991)
 - [7] O. Schönborn and R. Desai, Report No. cond-mat/9804237
 - [8] B. J. Kim, P. Minnhagen and P. Olsson, Report. No. cond-mat/9806231
 - [9] F. Liu, G. Mazenko, *Phys. Rev. B* **45**, 6989 (1992); **B 46**, 5963 (1992)
 - [10] O. Schöborn, R. Desai, Report No. cond-mat/9804237
 - [11] B. J. Kim, P. Minnhagen, P. Olsson, Report No. cond-mat/9806231
 - [12] A. C. Davis and R. Brandenberger, *Formation and Interactions of Topological defects*, NATO ASI, Ser. B. Vol. 349 (Plenum, New York, 1995)
 - [13] A. J. Bray, *Adv. Phys.* **43**, 357 (1994)
 - [14] Y. S. Duan, Y. Jiang and T. Xu, *Phys. Lett. A* **252**, 307 (1999)
 - [15] Y. S. Duan and X. H. Meng, *J. Math. Phys.* **34**, 1549 (1993)
 - [16] Y. S. Duan, and X. G. Lee, *Helv. Phys. Acta* **68**, 513 (1995)
 - [17] Y. S. Duan, S. Li and G. H. Yang, *Nucl. Phys.* **B514**, 705 (1998)
 - [18] Y. S. Duan and S. L. Zhang, *Int. J. Eng. Sci.* **28**, 689 (1990); **29**, 153 (1991); **29**, 1593 (1991); **30**, 153 (1992); Y. S. Duan and Z. P. Duan, *Int. J. Eng. Sci.* **24**, 531 (1986)
 - [19] Y. S. Duan, H. Zhang and S. Li, *Phys. Rev.* **B58** 125 (1998); Y. S. Duan, H. Zhang and L. B. Fu, *Phys. Rev.* **E59**, 528 (1999)

- [20] N. D. Mermin, *Rev. Mod. Phys.* **51**, 591 (1979)
- [21] M. Kleman, **Points, lines and walls, in liquid crystals, magnetic systems and various ordered media** (Wiley, New York, 1983)
- [22] Y. Jiang and Y. S. Duan, *Jour. Math. Phys.*, accepted (2000); Y. S. Duan and Y. Jiang, Report No. hep-th/9809011
- [23] I. M. Gel'fand and G. E. Shilov, *Generalized Functions*, Vol. 1, (Academic Press, New York, 1964); D. S. Jones, *Generalized Functions*, (McGraw-Hill, New York, 1966); E. M. De Jager, *Mathematics applied to physics* (E. Roubine, ed., Springer-Verlag, New York, 1970); J. A. Schouten, *Tensor Analysis for Physicists* (Oxford, Clarendon Press, 1951)
- [24] É. Goursat, *A Course in Mathematical Analysis* Vol.I, translated by Earle Raymond Hedrick, 1904
- [25] G. F. Mazenko, Report No. cond-mat/9808223
- [26] B. A. Dubrovin et. al., *Modern Geometry Method and Application*, Part II, Springer-Verlag New York Inc. 1985; Y. Choquet-Bruhat et. al, *Analysis, Manifolds and Physics*, North-Holland Publishing Company, Amsterdam, 1977
- [27] J. W. Milnor, *Topology, From the Differential Viewpoint*, The University Press of Virginia Charlottesville, 1965; S. Lefschetz, Introduction to Topology, Princeton University Press, Princeton, New Jersey, 1949
- [28] B. I. Halperin, in *Physics of Defects*, edited by R. Balian *et al.* (North-Holland, Amsterdam, 1981)
- [29] M. Kubiček and M. Marek, *Computational Methods in Bifurcation Theory and Dissipative Structures* (Springer-Verlag, New York Inc., 1983)

FIGURES' CAPTIONS

Fig. 1. (a) The creation of two topological defects. (b) Two topological defects annihilate in collision at the limit point.

Fig. 2. Two topological defects collide with different directions of motion at the bifurcation point.

Fig. 3. Topological defects have the same direction of motion. (a) Two topological defects tangentially collide at the bifurcation point. (b) Two topological defects merge into one topological defect at the bifurcation point. (c) One topological defect splits into two topological defects at the bifurcation point.

Fig. 4. (a) One topological defect splits into three topological defects at the bifurcation point. (b) Three topological defects merge into one topological defect at the bifurcation point.

Fig. 5. This case is similar to Fig. 4. (a) Three topological defects merge into one topological defect at the bifurcation point. (b) One topological defect splits into three topological defects at the bifurcation point.




Fig. 1(a)




Fig. 1(b)




Fig. 2




Fig. 3(a)




Fig. 3(b)




Fig. 3(c)




Fig. 4(a)




Fig. 4(b)




Fig. 5(a)




Fig. 5(b)